

Noether's second theorem in a general setting. Reducible gauge theories

D.Bashkirov¹, G.Giachetta², L.Mangiarotti², G.Sardanashvily¹

¹ Department of Theoretical Physics, Moscow State University, 117234 Moscow, Russia

² Department of Mathematics and Informatics, University of Camerino, 62032 Camerino (MC), Italy

Abstract

We prove Noether's direct and inverse second theorems for Lagrangian systems on fiber bundles in the case of gauge symmetries depending on derivatives of dynamic variables and parameters of an arbitrary order. The appropriate notions of a reducible gauge symmetry and Noether identity are formulated, and their equivalence by means a certain intertwining operator is proved.

Mathematical subject classification: Primary 70S05, Secondary 53C80, 58A20

1 Introduction

Different variants of Noether's second theorem state that, if a Lagrangian admits symmetries depending on parameters, its variational derivatives obey certain relations, called the Noether identity. In a rather general setting, this theorem has been formulated in [8]. Gauge symmetries and Noether identities need not be independent, and one speaks on N -stage reducible gauge symmetries and Noether identities. The notion of a reducible Noether identity has come from that of a reducible constraint [7], but it involves differential operators. Note that the conventional Batalin–Vilkovisky (BV) quantization of a classical gauge system necessarily starts with studying an hierarchy of its gauge symmetries and Noether identities in order to define the multiplet of ghosts and antifields, and to construct the so-called gauge-fixed Lagrangian [2, 13]. It should be also emphasized that, if a gauge symmetry is reducible, the components of a Noether current in classical field theory and the Ward identities in quantum field theory fail to be independent.

We present Noether's second theorem and its inverse (Theorem 4.2) for Lagrangian systems on a fiber bundle $Y \rightarrow X$ in the case of gauge symmetries depending on derivatives of dynamic variables and parameters of an arbitrary order. Bearing in mind the extension of the BV quantization scheme to an arbitrary base manifold [3, 12], we pay the particular attention to global aspects of Noether's second theorem. For this purpose, we consider Lagrangian formalism on the composite fiber bundle $E \rightarrow Y \rightarrow X$, where $E \rightarrow Y$ is a vector bundle of gauge parameters. Accordingly, a gauge symmetry is represented by a linear differential operator v on E taking its values in the vertical tangent bundle VY of $Y \rightarrow X$.

The Noether identity for a Lagrangian L is defined by a differential operator Δ on the fiber bundle (2.12) which takes its values in the density-dual

$$\overline{E}^* = E^* \otimes_Y^n T^*X \quad (1.1)$$

of $E \rightarrow Y$ and whose kernel contains the image of the Euler–Lagrange operator δL of L , i.e.,

$$\Delta \circ \delta L = 0. \quad (1.2)$$

Expressed in these terms, Noether’s second theorem and its inverse follow at once from the first variational formula (Proposition 3.1) and the properties of differential operators on dual fiber bundles (Theorem 8.1). Namely, there exists the intertwining operator $\eta(v) = \Delta$ (8.4), $\eta(\Delta) = v$ (8.5) such that

$$\eta(\eta(v)) = v, \quad \eta(\eta(\Delta)) = \Delta, \quad (1.3)$$

$$\eta(v \circ v') = \eta(v') \circ \eta(v), \quad \eta(\Delta' \circ \Delta) = \eta(\Delta) \circ \eta(\Delta'). \quad (1.4)$$

The appropriate notions of a reducible Noether identity and gauge symmetry are formulated, and their equivalence with respect to the intertwining operator η is proved (Theorem 5.3).

The following two examples aim to illustrate our exposition: (i) the gauge theory of principal connections for which gauge transformations need not be vertical, e.g., the topological gauge theory with the global Chern–Simons Lagrangian and the Yang–Mills gauge theory with a dynamic metric field, (ii) a gauge system of skew symmetric tensor fields with a reducible gauge symmetry, e.g., the topological BF theory.

2 Lagrangian formalism on fiber bundles

Lagrangian formalism on a fiber bundle $Y \rightarrow X$ is phrased in terms of the following graded differential algebra (henceforth GDA) [1, 10, 12, 15].

The finite order jet manifolds of $Y \rightarrow X$ form an inverse system

$$X \xleftarrow{\pi} Y \xleftarrow{\pi_0^1} J^1 Y \xleftarrow{\dots} J^{r-1} Y \xleftarrow{\pi_{r-1}^r} J^r Y \xleftarrow{\dots}. \quad (2.1)$$

In the sequel, the index $r = 0$ stands for Y . Accordingly, we have the direct system

$$\mathcal{O}^* X \xrightarrow{\pi^*} \mathcal{O}^* Y \xrightarrow{\pi_0^{1*}} \mathcal{O}_1^* Y \longrightarrow \dots \mathcal{O}_{r-1}^* Y \xrightarrow{\pi_{r-1}^{r*}} \mathcal{O}_r^* Y \longrightarrow \dots \quad (2.2)$$

of GDAs $\mathcal{O}_r^* Y$ of exterior forms on jet manifolds $J^r Y$ with respect to the pull-back monomorphisms π_{r-1}^{r*} . Its direct limit $\mathcal{O}_\infty^*[Y]$ is a GDA consisting of all exterior forms on finite order jet manifolds modulo the pull-back identification.

The projective limit $(J^\infty Y, \pi_r^\infty : J^\infty Y \rightarrow J^r Y)$ of the inverse system (2.1) is a Fréchet manifold. A bundle atlas $\{(U_Y; x^\lambda, y^i)\}$ of $Y \rightarrow X$ yields the coordinate atlas

$$\{((\pi_0^\infty)^{-1}(U_Y); x^\lambda, y_\Lambda^i)\}, \quad y_{\lambda+\Lambda}^i = \frac{\partial x^\mu}{\partial x'^\lambda} d_\mu y_\Lambda^i, \quad 0 \leq |\Lambda|, \quad (2.3)$$

of $J^\infty Y$, where $\Lambda = (\lambda_k \dots \lambda_1)$ is a symmetric multi-index, $\lambda + \Lambda = (\lambda \lambda_k \dots \lambda_1)$, and

$$d_\lambda = \partial_\lambda + \sum_{0 \leq |\Lambda|} y_{\lambda+\Lambda}^i \partial_i^\Lambda, \quad d_\Lambda = d_{\lambda_r} \circ \dots \circ d_{\lambda_1} \quad (2.4)$$

are total derivatives. There is the GDA epimorphism $\mathcal{O}_\infty^*[Y] \rightarrow \mathcal{O}_\infty^*[U_Y]$ obtained as the restriction of $\mathcal{O}_\infty^* Y$ to the chart (2.3). Then $\mathcal{O}_\infty^*[Y]$ can be written in a coordinate form where the horizontal one-forms $\{dx^\lambda\}$ and the contact one-forms $\{\theta_\Lambda^i = dy_\Lambda^i - y_{\lambda+\Lambda}^i dx^\lambda\}$ are generating elements of the $\mathcal{O}_\infty^0[U_Y]$ -algebra $\mathcal{O}_\infty^*[U_Y]$. Though $J^\infty Y$ is not a smooth manifold, the coordinate transformations of elements of $\mathcal{O}_\infty^*[Y]$ are smooth since they are exterior forms on finite order jet manifolds.

There is the canonical decomposition $\mathcal{O}_\infty^*[Y] = \oplus \mathcal{O}_\infty^{k,m}[Y]$ of $\mathcal{O}_\infty^*[Y]$ into $\mathcal{O}_\infty^0[Y]$ -modules $\mathcal{O}_\infty^{k,m}[Y]$ of k -contact and m -horizontal forms together with the corresponding projectors

$$h_k : \mathcal{O}_\infty^*[Y] \rightarrow \mathcal{O}_\infty^{k,*}[Y], \quad h^m : \mathcal{O}_\infty^*[Y] \rightarrow \mathcal{O}_\infty^{*,m}[Y].$$

Accordingly, the exterior differential on $\mathcal{O}_\infty^*[Y]$ is split into the sum $d = d_H + d_V$ of the nilpotent total and vertical differentials

$$d_H(\phi) = dx^\lambda \wedge d_\lambda(\phi), \quad d_V(\phi) = \theta_\Lambda^i \wedge \partial_i^\Lambda \phi, \quad \phi \in \mathcal{O}_\infty^*[Y].$$

In particular, any finite order Lagrangian on a fiber bundle $Y \rightarrow X$ is a density

$$L = \mathcal{L}\omega \in \mathcal{O}_\infty^{0,n}[Y], \quad \omega = dx^1 \wedge \dots \wedge dx^n, \quad n = \dim X. \quad (2.5)$$

In the framework of Lagrangian formalism, we deal with differential operators of the following type. Let

$$W \rightarrow Y \rightarrow X, \quad Z \rightarrow Y \rightarrow X$$

be composite bundles, including $W = Y$, and let $Z \rightarrow Y$ be a vector bundle. By a k -order differential operator on $W \rightarrow X$ taking its values in $Z \rightarrow X$ is throughout meant a bundle morphism

$$\Delta : J^k W \xrightarrow{Y} Z. \quad (2.6)$$

Its kernel $\text{Ker } \Delta$ is defined as the inverse image of the canonical zero section of $Z \rightarrow Y$. In an equivalent way, the differential operator (2.6) is represented by a section Δ of the vector

bundle $J^k W \times_Y Z \rightarrow J^k W$. Given bundle coordinates (x^λ, y^i, w^r) on W and (x^λ, y^i, z^A) on Z with respect to the fiber basis $\{e_A\}$ for $Z \rightarrow Y$, this section reads

$$\Delta = \Delta^A(x^\lambda, y_\Lambda^i, w_\Lambda^B) e_A, \quad 0 \leq |\Lambda| \leq k. \quad (2.7)$$

Then the differential operator (2.6) is also represented by an element

$$\Delta = \Delta^A(x^\lambda, y_\Lambda^i, w_\Lambda^B) z_A \in \mathcal{O}_\infty^0[W \times Z^*] \quad (2.8)$$

of the GDA $\mathcal{O}_\infty^*[W \times_X Z^*]$, where $Z^* \rightarrow Y$ is the dual of $Z \rightarrow Y$ with coordinates (x^λ, y^i, z_A) .

If $W \rightarrow Y$ is a vector bundle, a differential operator Δ (2.6) on the composite bundle $W \rightarrow Y \rightarrow X$ is said to be linear if it is linear on the fibers of the vector bundle $J^k W \rightarrow J^k Y$. In this case, its representations (2.7) and (2.8) take the form

$$\Delta = \sum_{0 \leq |\Xi| \leq k} \Delta_r^{A, \Xi}(x^\lambda, y_\Lambda^i) w_\Xi^r e_A, \quad 0 \leq |\Lambda| \leq k, \quad (2.9)$$

$$\Delta = \sum_{0 \leq |\Xi| \leq k} \Delta_r^{A, \Xi}(x^\lambda, y_\Lambda^i) w_\Xi^r z_A, \quad 0 \leq |\Lambda| \leq k. \quad (2.10)$$

In particular, every Lagrangian L (2.5) defines the Euler–Lagrange operator

$$\delta L = \sum_{0 \leq |\Lambda|} (-1)^{|\Lambda|} d_\Lambda(\partial_i^\Lambda \mathcal{L}) dy^i \wedge \omega \quad (2.11)$$

on Y taking the values in the vector bundle

$$V^* Y \otimes_Y \bigwedge_X^n T^* X \rightarrow Y. \quad (2.12)$$

It is represented by the exterior form

$$\delta L = \mathcal{E}_i \theta^i \wedge \omega = \sum_{0 \leq |\Lambda|} (-1)^{|\Lambda|} d_\Lambda(\partial_i^\Lambda \mathcal{L}) \theta^i \wedge \omega \in \mathcal{O}_\infty^{1, n}[Y], \quad (2.13)$$

where

$$\delta \phi = \sum_{0 \leq |\Lambda|} (-1)^{|\Lambda|} \theta^i \wedge [d_\Lambda(\partial_i^\Lambda) d\phi], \quad \phi \in \mathcal{O}_\infty^{*, n}[Y], \quad (2.14)$$

is the variational operator acting on $\mathcal{O}_\infty^{*, n}[Y]$ so that $\delta \circ d_H = 0$ and $\delta \circ \delta = 0$. There is the canonical decomposition

$$dL = \delta L - d_H \Xi, \quad (2.15)$$

where $\Xi_L = L + \Xi$ is a Lepagean equivalent of L . It reads

$$\begin{aligned} \Xi_L &= L + \sum_{s=0} F_i^{\lambda \nu_s \dots \nu_1} \theta_{\nu_s \dots \nu_1}^i \wedge \omega_\lambda, \\ F_i^{\nu_k \dots \nu_1} &= \partial_i^{\nu_k \dots \nu_1} \mathcal{L} - d_\lambda F_i^{\lambda \nu_k \dots \nu_1} + h_i^{\nu_k \dots \nu_1}, \quad \omega_\lambda = \partial_\lambda \rfloor \omega, \end{aligned}$$

where functions h obey the relations $h_i^\nu = 0$, $h_i^{(\nu_k \nu_{k-1} \dots \nu_1)} = 0$ [14].

Remark 2.1. Given a Lagrangian L and its Euler-Lagrange operator δL (2.13), we further abbreviate $A \approx 0$ with an equality which holds on-shell. This means that A is an element of a module over the ideal I_L of the ring $\mathcal{O}_\infty^0[Y]$ which is locally generated by the variational derivatives $\mathcal{E}_i$ and their total derivatives $d_\Lambda \mathcal{E}_i$. One says that I_L is a differential ideal because, if a local functions f belongs to I_L , then every total derivative $d_\Lambda f$ does as well.

Remark 2.2. We will use the relations

$$\sum_{0 \leq |\Lambda| \leq k} B^\Lambda d_\Lambda A' = \sum_{0 \leq |\Lambda| \leq k} (-1)^{|\Lambda|} d_\Lambda (B^\Lambda) A' + d_H \sigma, \quad (2.16)$$

$$\sum_{0 \leq |\Lambda| \leq k} (-1)^{|\Lambda|} d_\Lambda (B^\Lambda A) = \sum_{0 \leq |\Lambda| \leq k} \eta(B)^\Lambda d_\Lambda A, \quad (2.17)$$

$$\eta(B)^\Lambda = \sum_{0 \leq |\Sigma| \leq k - |\Lambda|} (-1)^{|\Sigma + \Lambda|} C_{|\Sigma + \Lambda|}^{|\Sigma|} d_\Sigma B^{\Sigma + \Lambda}, \quad C_b^a = \frac{b!}{a!(b-a)!}, \quad (2.18)$$

for arbitrary exterior forms $A' \in \mathcal{O}_\infty^{*,n}[Q]$, $A \in \mathcal{O}_\infty^*[Q]$ and local functions $B^\Lambda \in \mathcal{O}_\infty^0[Q]$ on jet manifolds of a fiber bundle $Q \rightarrow X$. Since $\sum_{a=0}^k (-1)^a C_k^a = 0$ for $k > 0$, it is easily verified that

$$(\eta \circ \eta)(B)^\Lambda = B^\Lambda. \quad (2.19)$$

3 Gauge symmetries in a general setting

Let $\mathfrak{d}\mathcal{O}_\infty^0 Y$ be the $\mathcal{O}_\infty^0[Y]$ -module of derivations of the $\mathbb{R}$ -algebra $\mathcal{O}_\infty^0[Y]$. Any $\vartheta \in \mathfrak{d}\mathcal{O}_\infty^0 Y$ yields a graded derivation (the interior product) $\vartheta \rfloor \phi$ of the GDA $\mathcal{O}_\infty^*[Y]$ given by the relations

$$\begin{aligned} \vartheta \rfloor df &= \vartheta(f), & f &\in \mathcal{O}_\infty^0[Y], \\ \vartheta \rfloor (\phi \wedge \sigma) &= (\vartheta \rfloor \phi) \wedge \sigma + (-1)^{|\phi|} \phi \wedge (\vartheta \rfloor \sigma), & \phi, \sigma &\in \mathcal{O}_\infty^*[Y], \end{aligned}$$

and a derivation $\mathbf{L}_\vartheta$ (the Lie derivative) which satisfies the conditions

$$\mathbf{L}_\vartheta \phi = \vartheta \rfloor d\phi + d(\vartheta \rfloor \phi), \quad \phi \in \mathcal{O}_\infty^*[Y], \quad (3.1)$$

$$\mathbf{L}_\vartheta(\phi \wedge \phi') = \mathbf{L}_\vartheta(\phi) \wedge \phi' + \phi \wedge \mathbf{L}_\vartheta(\phi'), \quad (3.2)$$

$$\mathbf{L}_\vartheta(d_H \phi) = d_H(\mathbf{L}_\vartheta \phi). \quad (3.3)$$

Relative to an atlas (2.3), a derivation $\vartheta \in \mathfrak{d}\mathcal{O}_\infty^0 Y$ reads

$$\vartheta = \vartheta^\lambda \partial_\lambda + \vartheta^i \partial_i + \sum_{|\Lambda| > 0} \vartheta_\Lambda^i \partial_i^\Lambda, \quad (3.4)$$

where the tuple of derivations $\{\partial_\lambda, \partial_i^\Lambda\}$ is defined as the dual of the set $\{dx^\lambda, dy_\lambda^i\}$ of generating elements for the $\mathcal{O}_\infty^0[Y]$ -algebra $\mathcal{O}_\infty^*[Y]$ with respect to the interior product $\rfloor$ [12].

A derivation ϑ (3.4) is called contact if the Lie derivative $\mathbf{L}_\vartheta$ (3.1) preserves the contact ideal of the GDA $\mathcal{O}_\infty^*[Y]$ generated by contact forms. A derivation ϑ (3.4) is contact iff

$$\vartheta_\Lambda^i = d_\Lambda(\vartheta^i - y_\mu^i \vartheta^\mu) + y_{\mu+\Lambda}^i \vartheta^\mu, \quad 0 < |\Lambda|. \quad (3.5)$$

Any contact derivation admits the canonical horizontal splitting

$$\vartheta = \vartheta_H + \vartheta_V = \vartheta^\lambda d_\lambda + (v^i \partial_i + \sum_{0 < |\Lambda|} d_\Lambda v^i \partial_i^\Lambda), \quad v^i = \vartheta^i - y_\mu^i \vartheta^\mu. \quad (3.6)$$

Its vertical part ϑ_V is completely determined by the first summand

$$v = v^i(x^\lambda, y_\Lambda^i) \partial_i, \quad 0 \leq |\Lambda| \leq k. \quad (3.7)$$

This is a section of the pull-back $VY \times_Y J^k Y \rightarrow J^k Y$ of the vertical tangent bundle $VY \rightarrow Y$ onto $J^k Y$ [6] and, thus, it is a k -order VY -valued differential operator on Y . One calls v (3.7) a generalized vector field on Y .

Proposition 3.1. *It follows from the splitting (2.15) that the Lie derivative of a Lagrangian L (2.5) along a contact derivation ϑ (3.9) fulfills the first variational formula*

$$\mathbf{L}_\vartheta L = v \rfloor \delta L + d_H(h_0(\vartheta \rfloor \Xi_L)) + \mathcal{L}d_V(\vartheta_H \rfloor \omega), \quad (3.8)$$

where Ξ_L is a Lepagean equivalent of L [12].

A contact derivation ϑ (3.9) is called variational if the Lie derivative (3.8) is d_H -exact, i.e., $\mathbf{L}_\vartheta L = d_H \sigma$, $\sigma \in \mathcal{O}_\infty^{0,n-1}[Y]$. A glance at the expression (3.8) shows that: (i) a contact derivation ϑ is variational only if it is projectable onto X , (ii) ϑ is variational iff its vertical part ϑ_V is well, (iii) it is variational if $v \rfloor \delta L$ is d_H -exact. By virtue of item (ii), we can restrict our consideration to vertical contact derivations

$$\vartheta = \sum_{0 \leq |\Lambda|} d_\Lambda v^i \partial_i^\Lambda. \quad (3.9)$$

A generalized vector field v (3.7) is called a variational symmetry of a Lagrangian L if it generates a variational vertical contact derivation (3.9).

A Lagrangian system on a fiber bundle $Y \rightarrow X$ is said to be a gauge theory if its Lagrangian L admits a family of variational symmetries parameterized by elements of a vector bundle $E \rightarrow Y$ as follows.

Let $E \rightarrow Y$ be a vector bundle coordinated by (x^λ, y^i, ξ^r) . Given a Lagrangian L on Y , let us consider its pull-back, say again L , onto E . Let ϑ_E be a contact derivation of the $\mathbb{R}$ -algebra $\mathcal{O}_\infty^0[E]$, whose restriction

$$\vartheta = \vartheta_E|_{\mathcal{O}_\infty^0 Y} = \sum_{0 \leq |\Lambda|} d_\Lambda v^i \partial_i^\Lambda \quad (3.10)$$

to $\mathcal{O}_\infty^0[Y] \subset \mathcal{O}_\infty^0[E]$ is linear in coordinates ξ_Ξ^r . It is determined by a generalized vector field v_E on E whose canonical projection

$$v : J^k E \xrightarrow{v_E} VE \rightarrow E \times_Y VY$$

(see the exact sequence (3.12) below) is a linear VY -valued differential operator

$$v = \sum_{0 \leq |\Xi| \leq m} v_r^{i, \Xi}(x^\lambda, y_\Sigma^i) \xi_\Xi^r \partial_i \quad (3.11)$$

on $E \rightarrow Y \rightarrow X$. Let ϑ_E be a variational symmetry of a Lagrangian L on E , i.e.,

$$\mathbf{L}_{\vartheta_E} L = d_H \sigma.$$

Then one says that v (3.11) is a gauge symmetry of a Lagrangian L .

Remark 3.1. Note that any generalized vector field v (3.11) gives rise to a generalized vector field v_E on E and, thus, defines a contact derivation ϑ_E of $\mathcal{O}_\infty^0[E]$. Indeed, let us consider the exact sequence of vector bundles

$$0 \rightarrow V_Y E \rightarrow VE \rightarrow E \times_Y VY \rightarrow 0, \quad (3.12)$$

where $V_Y E$ is the vertical tangent bundle of $E \rightarrow Y$. Its splitting Γ lifts v to the generalized vector field $v_E = \Gamma \circ v$ on E such that the Lie derivative

$$\mathbf{L}_{\vartheta_E} L = v \rfloor \delta L + d_H(\vartheta \rfloor \Xi_L) \quad (3.13)$$

depends only on v , but not a lift Γ .

Remark 3.2. If v (3.11) is a gauge symmetry, we obtain from the first variational formula (3.13) the weak conservation law

$$0 \approx d_H(\vartheta \rfloor \Xi_L - \sigma), \quad (3.14)$$

where

$$J = \vartheta \rfloor \Xi_L = \sum_{0 \leq |\Lambda|} J_r^{\lambda, \Lambda} \xi_\Lambda^r \omega_\lambda, \quad (3.15)$$

is a Noether current.

4 Noether's second theorem

Let us start with the notion of a Noether identity.

Definition 4.1. *Given a Lagrangian L (2.5) and its Euler–Lagrange operator δL (2.13), let $E \rightarrow Y$ be a vector bundle and Δ a linear differential operator of order $0 \leq m$ on the composite bundle (2.12) with the values in the density-dual $\overline{E}^*$ (1.1) of E which obeys the condition (1.2). This condition is called the Noether identity, and Δ is the Noether operator.*

Given bundle coordinates $(x^\lambda, y^i, \overline{y}_i)$ on the fiber bundle (2.12) and (x^λ, y^i, ξ^r) on E , the Noether operator Δ is represented by the density

$$\Delta = \Delta_r \xi^r \omega = \sum_{0 \leq |\Lambda| \leq m} \Delta_r^{i,\Lambda} (x^\lambda, y_\Sigma^j) \overline{y}_{\Lambda i} \xi^r \omega \in \mathcal{O}_\infty^{0,n}[E \times_Y V^* Y], \quad 0 \leq |\Sigma| \leq m. \quad (4.1)$$

Then the Noether identity (1.2) takes the coordinate form

$$\sum_{0 \leq |\Lambda| \leq m} \Delta_r^{i,\Lambda} d_\Lambda \mathcal{E}_i \xi^r \omega = 0. \quad (4.2)$$

Theorem 4.2. *If a Lagrangian L (2.5) admits a gauge symmetry v (3.11), its Euler–Lagrange operator obeys the Noether identity (4.2) where the Noether operator (4.1) is*

$$\begin{aligned} \Delta = \eta(v) &= \sum_{0 \leq |\Sigma| \leq m} (-1)^{|\Sigma|} d_\Sigma (v_r^{i,\Sigma} \overline{y}_i) \xi^r \omega = \sum_{0 \leq |\Lambda| \leq m} \eta(v)_r^{i,\Lambda} \overline{y}_{\Lambda i} \xi^r \omega, \\ \eta(v)_r^{i,\Lambda} &= \sum_{0 \leq |\Sigma| \leq m-|\Lambda|} (-1)^{|\Sigma+\Lambda|} C_{|\Sigma+\Lambda|}^{|\Sigma|} d_\Sigma v_r^{i,\Sigma+\Lambda}. \end{aligned} \quad (4.3)$$

Conversely, if the Euler–Lagrange operator of a Lagrangian L obeys the Noether identity (4.2), this Lagrangian admits a gauge symmetry v (3.11) where

$$v = \eta(\Delta) = \sum_{0 \leq |\Sigma| \leq m} (-1)^{|\Sigma|} d_\Sigma (\Delta_r^{i,\Sigma} \xi^r) \partial_i = \sum_{0 \leq |\Lambda| \leq m} \eta(\Delta)_r^{i,\Lambda} \xi_\Lambda^r \partial_i, \quad (4.4)$$

$$\eta(\Delta)_r^{i,\Lambda} = \sum_{0 \leq |\Sigma| \leq m-|\Lambda|} (-1)^{|\Sigma+\Lambda|} C_{|\Sigma+\Lambda|}^{|\Sigma|} d_\Sigma \Delta_r^{i,\Sigma+\Lambda}. \quad (4.5)$$

Proof. Given an operator v (3.11), the operator $\Delta = \eta(v)$ (4.3) is defined in accordance with Theorem 8.1 in Appendix. Since the density

$$v \rfloor \delta L = v^i \mathcal{E}_i \omega = \sum_{0 \leq |\Xi| \leq m} v_r^{i,\Xi} \xi_\Xi^r \mathcal{E}_i \omega$$

is d_H -exact, the Noether identity

$$\delta(v \rfloor \delta L) = \eta(v) \circ \delta L = 0$$

holds. Conversely, any operator Δ (4.1) defines the generalized vector field $v = \eta(\Delta)$ (4.4). Due to the Noether identity (4.2), we obtain

$$\begin{aligned} 0 &= \sum_{0 \leq |\Lambda| \leq m} \xi^r \Delta_r^{i,\Lambda} d_\Lambda \mathcal{E}_i \omega = \sum_{0 \leq |\Lambda| \leq m} (-1)^{|\Lambda|} d_\Lambda (\xi^r \Delta_r^{i,\Lambda}) \mathcal{E}_i \omega + d_H \sigma = \\ &\quad \sum_{0 \leq |\Xi| \leq m} v_r^{i,\Xi} \xi_\Xi^r \mathcal{E}_i \omega + d_H \sigma = v \rfloor \delta L + d_H \sigma, \end{aligned}$$

i.e., v is a gauge symmetry of L . $\square$

By virtue of the relations (1.3), there is one-to-one correspondence between gauge symmetries of a Lagrangian L and the Noether identities for δL .

Example 4.1. If a gauge symmetry

$$v = (v_r^i \xi^r + v_r^{i,\mu} \xi_\mu^r) \partial_i \quad (4.6)$$

is of first jet order in parameters, the corresponding Noether operator (4.3) and Noether identity take the form

$$\Delta = [(v_r^i - d_\mu v_r^{i,\mu}) \bar{y}_i - v_r^{i,\mu} \bar{y}_{\mu i}] \xi^r \omega, \quad (4.7)$$

$$[v_r^i \mathcal{E}_i - d_\mu (v_r^{i,\mu} \mathcal{E}_i)] \xi^r \omega = 0. \quad (4.8)$$

Any Lagrangian L has gauge symmetries. In particular, there always exist trivial gauge symmetries

$$v = \sum_\Lambda \eta(M)_r^{i,\Lambda} \xi_\Lambda^r, \quad M_r^{i,\Lambda} = \sum_\Sigma T_r^{i,j,\Lambda,\Sigma} d_\Sigma \mathcal{E}_j, \quad T_r^{j,i,\Lambda,\Sigma} = -T_r^{i,j,\Sigma,\Lambda},$$

corresponding to the trivial Noether identity

$$\sum_{\Sigma, \Lambda} T_r^{j,i,\Lambda,\Sigma} d_\Sigma \mathcal{E}_j d_\Lambda \mathcal{E}_i = 0.$$

Furthermore, given a gauge symmetry v (3.11), let $E' \rightarrow Y$ be a vector bundle and h a linear differential operator on some composite bundle $E' \rightarrow Y \rightarrow X$, coordinated by (x^λ, y^i, ξ'^s) , with values in the vector bundle $E \rightarrow Y$. Then the composition

$$v' = v \circ h = v_s^{i,\Lambda} \xi_\Lambda'^s \partial_i, \quad v_s^{i,\Lambda} = \sum_{\Xi + \Xi' = \Lambda} \sum_{0 \leq |\Sigma| \leq m - |\Xi|} v_r^{i,\Xi + \Sigma} d_\Sigma h_s^{r,\Xi'},$$

is a variational symmetry of the pull-back of a Lagrangian L onto E' , i.e., a gauge symmetry. In view of this ambiguity, we agree to say that a gauge symmetry v (3.11) of a Lagrangian L is complete if any different gauge symmetry v'_0 of L factorizes through v as

$$v' = v \circ h + T, \quad T \approx 0.$$

A complete gauge symmetry always exists, but the vector bundle of its parameters need not be finite-dimensional.

Accordingly, given the Noether operator (4.1), let H be a linear differential operator on $\overline{E}^* \rightarrow Y \rightarrow X$ with values in the density-dual $\overline{E}'^*$ (1.1) of some vector bundle $E' \rightarrow Y$. Then the composition $\Delta' = H \circ \Delta$ is also a Noether operator. We agree to call the Noether operator (4.1) complete if a different Noether operator Δ' factors through Δ as

$$\Delta' = H \circ \Delta + F, \quad F \approx 0.$$

Proposition 4.3. *A gauge symmetry v of a Lagrangian L is complete iff the associated Noether operator is also.*

Proof. The proof follows at once from Proposition 8.2 in Appendix. Given a gauge symmetry v of L , let v' be a different gauge symmetry. If $\eta(v)$ is a complete Noether operator, then

$$\eta(v') = H \circ \eta(v) + F, \quad F \approx 0,$$

and, by virtue of the relations (1.4), we have

$$v' = v \circ \eta(H) + \eta(F),$$

where $\eta(F) \approx 0$ because I_L is a differential ideal. The converse is similarly proved. $\square$

5 Reducible gauge theories

Let us extend Noether's second theorem to the analysis of reducible gauge systems.

Definition 5.1. *A complete Noether operator $\Delta \not\approx 0$ (4.1) and the corresponding Noether identity (1.2) are said to be N -stage reducible ($N = 0, 1, \dots$) if there exist vector bundles $E_k \rightarrow Y$ and differential operators Δ_k , $k = 0, \dots, N$, such that:*

(i) Δ_k is a linear differential operator on the density-dual $\overline{E}_{k-1}^* \rightarrow Y \rightarrow X$ of $E_{k-1} \rightarrow Y$ with values in the density-dual $\overline{E}_k^*$ of E_k , where $E_{-1} = E$;

(ii) $\Delta_k \not\approx 0$ for all $k = 0, \dots, N$;

(iii) $\Delta_k \circ \Delta_{k-1} \approx 0$ for all $k = 0, \dots, N$, where $\Delta_{-1} = \Delta$;

(iv) if Δ'_k is another differential operator possessing these properties, then it factors through Δ_k on-shell.

In particular, a zero-stage reducible Noether operator is called reducible. In this case, given bundle coordinates $(x^\lambda, y^i, \bar{\xi}_r)$ on $\bar{E}^*$ and $(x^\lambda, y^i, \xi^{r_0})$ on E_0 , a differential operator Δ_0 reads

$$\Delta_0 = \sum_{0 \leq |\Xi| \leq m_0} \Delta_{r_0}^{r, \Xi} \bar{\xi}_{\Xi r} \xi^{r_0} \omega. \quad (5.1)$$

Then the reduction condition $\Delta_0 \circ \Delta \approx 0$ takes the coordinate form

$$\sum_{0 \leq |\Xi| \leq m_0} \Delta_{r_0}^{r, \Xi} d_{\Xi} \left(\sum_{0 \leq |\Lambda| \leq m} \Delta_r^{i, \Lambda} \bar{y}_{\Lambda i} \right) \xi^{r_0} \omega \approx 0, \quad (5.2)$$

i.e., the left hand-side of this expression takes the form

$$\sum_{0 \leq |\Sigma| \leq m_0 + m} M_{r_0}^{i, \Sigma} \bar{y}_{\Sigma i} \xi^{r_0} \omega,$$

where all the coefficients $M_{r_0}^{i, \Sigma}$ belong to the ideal I_L .

Definition 5.2. A complete gauge symmetry $v \not\approx 0$ (3.11) is said to be N -stage reducible if there exist vector bundles $E_k \rightarrow Y$ and differential operators v^k , $k = 0, \dots, N$, such that:

(i) v^k is a linear differential operator on the composite bundle $E_k \rightarrow Y \rightarrow X$ with values in the vector bundle $E_{k-1} \rightarrow Y$;

(ii) $v^k \not\approx 0$ for all $k = 0, \dots, N$;

(iii) $v^{k-1} \circ v^k \approx 0$ for all $k = 0, \dots, N$, where v^k , $k = -1$, stands for v ;

(iv) if v'^k is another differential operator possessing these properties, then v'^k factors through v^k on-shell.

Theorem 5.3. A gauge symmetry v is N -stage reducible iff the associated Noether identity is also.

Proof. The proof follows at once from Theorem 8.1 and Proposition 8.2 in Appendix. Let us put $\Delta_k = \eta(v^k)$, $k = 0, \dots, N$. If $v^k \approx 0$, then $\eta(v^k) \approx 0$ because I_L is a differential ideal. By the same reason, if v^{k-1} and v^k obey the reduction condition $v^{k-1} \circ v^k \approx 0$, then

$$\eta(v^{k-1} \circ v^k) = \eta(v^k) \circ \eta(v^{k-1}) \approx 0.$$

The converse is justified in a similar way. The equivalence of the conditions in items (iv) of Definitions 5.1 and 5.2 is proved similarly to that in Proposition 4.3. $\square$

Remark 5.1. Let a gauge symmetry v (3.11) be reducible. Given bundle coordinates $(x^\lambda, y^i, \xi^{r_0})$ on E_0 , the differential operator v^0 reads

$$v^0 = \sum_{0 \leq |\Lambda| \leq m_0} v_{r_0}^{r, \Lambda} \xi_\Lambda^{r_0} \partial_r,$$

and the reduction condition $v \circ v^0 \approx 0$ takes the form

$$\sum_{0 \leq |\Xi| \leq m} v_r^{i, \Xi} d_\Xi \left(\sum_{0 \leq |\Lambda| \leq m_0} v_{r_0}^{r, \Lambda} \xi_\Lambda^{r_0} \right) \partial_i \approx 0.$$

In particular, it follows that the Noether current J (3.15) vanishes on-shell if

$$\xi^r = \sum_{0 \leq |\Lambda| \leq m_0} v_{r_0}^{r, \Xi} \xi_\Lambda^{r_0}$$

and, consequently, its components $J_r^{\lambda, \Lambda}$ are not independent.

6 Example I

This example addresses the gauge model of principal connections on a principal bundle $P \rightarrow X$ with a structure Lie group G whose automorphisms need not be vertical. In a general setting, the gauge-natural prolongations of P and the associated natural-gauge bundles can be considered [6].

Principal connections on a principal bundle $P \rightarrow X$ are represented by sections of the quotient

$$C = J^1 P / G \rightarrow X, \tag{6.1}$$

called the bundle of principal connections. This is an affine bundle coordinated by (x^λ, a_λ^r) such that, given a section A of $C \rightarrow X$, its components $A_\lambda^r = a_\lambda^r \circ A$ are coefficients of the familiar local connection form (i.e., gauge potentials). We consider the GDA $\mathcal{O}_\infty^*[C]$.

Infinitesimal generators of one-parameter groups of automorphisms of a principal bundle P are G -invariant projectable vector fields on $P \rightarrow X$. They are associated to sections of the vector bundle $T_G P = TP/G \rightarrow X$. This bundle is endowed with the coordinates $(x^\lambda, \tau^\lambda = \dot{x}^\lambda, \xi^r)$ with respect to the fiber bases $\{\partial_\lambda, e_r\}$ for $T_G P$, where $\{e_r\}$ is the basis for the right Lie algebra $\mathfrak{g}$ of G such that $[e_p, e_q] = c_{pq}^r e_r$. If

$$u = u^\lambda \partial_\lambda + u^r e_r, \quad v = v^\lambda \partial_\lambda + v^r e_r, \tag{6.2}$$

are sections of $T_G P \rightarrow X$, their bracket reads

$$[u, v] = (u^\mu \partial_\mu v^\lambda - v^\mu \partial_\mu u^\lambda) \partial_\lambda + (u^\lambda \partial_\lambda v^r - v^\lambda \partial_\lambda u^r + c_{pq}^r u^p v^q) e_r.$$

Any section u of the vector bundle $T_GP \rightarrow X$ yields the vector field

$$u_C = u^\lambda \partial_\lambda + (c_{pq}^r a_\lambda^p u^q + \partial_\lambda u^r - a_\mu^r \partial_\lambda u^\mu) \partial_r^\lambda \quad (6.3)$$

on the bundle of principal connections C (6.1) [9].

In order to describe a gauge symmetry in this gauge model, let us consider the bundle product

$$E = C \times_X T_GP, \quad (6.4)$$

coordinated by $(x^\lambda, a_\lambda^r, \tau^\lambda, \xi^r)$. It can be provided with the generalized vector field

$$v_E = v = (c_{pq}^r a_\lambda^p \xi^q + \xi_\lambda^r - a_\mu^r \tau_\lambda^\mu - \tau^\mu a_{\mu\lambda}^r) \partial_r^\lambda. \quad (6.5)$$

With a subbundle $V_GP = VP/G \rightarrow X$ of the vector bundle T_GP coordinated by (x^λ, ξ^r) , we have the exact sequence of vector bundles

$$0 \rightarrow V_GP \longrightarrow T_GP \rightarrow TX \rightarrow 0.$$

The pull-back of this exact sequence via C admits the canonical splitting which takes the coordinate form

$$\tau^\lambda \partial_\lambda + \xi^r e_r = \tau^\lambda (\partial_\lambda + a_\lambda^r e_r) + (\xi^r - \tau^\lambda a_\lambda^r) e_r. \quad (6.6)$$

Due to this splitting, the generalized vector field (6.5) is brought into the form

$$v = (c_{pq}^r a_\lambda^p \xi'^q + \xi_\lambda'^r + \tau^\mu \mathcal{F}_{\lambda\mu}^r) \partial_r^\lambda, \quad \xi'^r = \xi^r - \tau^\lambda a_\lambda^r. \quad (6.7)$$

This generalized vector field is a gauge symmetry of the global Chern–Simons Lagrangian in gauge theory on a principal bundle with a structure semi-simple Lie group G over a three-dimensional base X . Given a section B of $C \rightarrow X$ (i.e., a background gauge potential), this Lagrangian reads

$$\begin{aligned} L = & \left[\frac{1}{2} a_{mn}^G \varepsilon^{\alpha\beta\gamma} a_\alpha^m (\mathcal{F}_{\beta\gamma}^n - \frac{1}{3} c_{pq}^n a_\beta^p a_\gamma^q) - \frac{1}{2} a_{mn}^G \varepsilon^{\alpha\beta\gamma} B_\alpha^m (F(B)_{\beta\gamma}^n \right. \\ & \left. - \frac{1}{3} c_{pq}^n B_\beta^p B_\gamma^q) - d_\alpha (a_{mn}^G \varepsilon^{\alpha\beta\gamma} a_\beta^m B_\gamma^n) \right] d^3x, \\ F(B)_{\lambda\mu}^r = & \partial_\lambda B_\mu^r - \partial_\mu B_\lambda^r + c_{pq}^r B_\lambda^p B_\mu^q, \quad \mathcal{F}_{\lambda\mu}^r = a_{\lambda\mu}^r - a_{\mu\lambda}^r + c_{pq}^r a_\lambda^p a_\mu^q, \end{aligned} \quad (6.8)$$

where a^G is the Killing form [5, 11]. Its first term is the well-known local Chern–Simons Lagrangian, the second one is a density on X , and the Lie derivative of the third term is d_H -exact due to the relation (3.3). The corresponding Noether identities (4.8) read

$$c_{pq}^r a_\lambda^p \mathcal{E}_r^\lambda - d_\lambda (\mathcal{E}_q^\lambda) = 0, \quad (6.9)$$

$$-a_{\mu\lambda}^r \mathcal{E}_r^\lambda + d_\lambda (a_\mu^r \mathcal{E}_r^\lambda) = 0. \quad (6.10)$$

The first one is the well-known Noether identity corresponding to the vertical gauge symmetry

$$v = (c_{pq}^r a_\lambda^p \xi^q + \xi_\lambda^r) \partial_r^\lambda.$$

The second Noether identity (6.10) is brought into the form

$$-a_\mu^r [c_{pq}^r a_\lambda^p \mathcal{E}_r^\lambda - d_\lambda(\mathcal{E}_q^\lambda)] + \mathcal{F}_{\lambda\mu}^r \mathcal{E}_r^\lambda = 0,$$

i.e., it is equivalent to the Noether identity $\mathcal{F}_{\lambda\mu}^r \mathcal{E}_r^\lambda = 0$, which also comes from the splitting (6.6) of the generalized vector field v . This Noether identity however is trivial since $\mathcal{F}_{\lambda\mu}^r = 0$ is the kernel of the Euler–Lagrange operator of the Chern–Simons Lagrangian (6.8).

In order to obtain a gauge symmetry of the Yang–Mills Lagrangian, one should complete the generalized vector field (6.5) with the term acting on a world metric.

Let LX the fiber bundle of linear frames in the tangent bundle TX of X . It is a principal bundle with the structure group $GL(n, \mathbb{R})$, $n = \dim X$, which admits reductions to its maximal compact subgroup $O(n)$. Global sections of the quotient bundle $\Sigma = LX/O(n)$ are Riemannian metrics on X . If X obeys the well-known topological conditions, pseudo-Riemannian metrics on X are similarly described. Being an open subbundle of the tensor bundle $\overset{2}{\vee} TX$, the fiber bundle Σ is provided with bundle coordinates $\sigma^{\mu\nu}$. It admits the canonical lift

$$u_\Sigma = u^\lambda \partial_\lambda + (\sigma^{\nu\beta} \partial_\nu u^\alpha + \sigma^{\alpha\nu} \partial_\nu u^\beta) \frac{\partial}{\partial \sigma^{\alpha\beta}} \quad (6.11)$$

of any vector field $u = u^\lambda \partial_\lambda$ on X . We describe the gauge system of principal connections and a dynamic metric field on the bundle product

$$E = C \times_X \Sigma \times_X T_G P, \quad (6.12)$$

coordinated by $(x^\lambda, a_\lambda^r, \sigma^{\alpha\beta}, \tau^\lambda, \xi^r)$. It can be provided with the generalized vector field

$$v = (c_{pq}^r a_\lambda^p \xi^q + \xi_\lambda^r - a_\mu^r \tau_\lambda^\mu - \tau^\mu a_{\mu\lambda}^r) \partial_r^\lambda + (\sigma^{\nu\beta} \tau_\nu^\alpha + \sigma^{\alpha\nu} \tau_\nu^\beta - \tau^\lambda \sigma_\lambda^{\alpha\beta}) \frac{\partial}{\partial \sigma^{\alpha\beta}}. \quad (6.13)$$

This is a gauge symmetry of the sum of the Yang–Mills Lagrangian $L_{\text{YM}}(\mathcal{F}_{\alpha\beta}^r, \sigma^{\mu\nu})$ and a Lagrangian of a metric field. The corresponding Noether identities read

$$\begin{aligned} c_{pq}^r a_\lambda^p \mathcal{E}_r^\lambda - d_\lambda(\mathcal{E}_q^\lambda) &= 0, \\ -a_{\mu\lambda}^r \mathcal{E}_r^\lambda + d_\lambda(a_\mu^r \mathcal{E}_r^\lambda) - \sigma_\mu^{\alpha\beta} \mathcal{E}_{\alpha\beta} - 2d_\nu(\sigma^{\nu\beta} \mathcal{E}_{\mu\beta}) &= 0. \end{aligned}$$

The first one is the Noether identity (6.9). Then the second identity is brought into the form

$$\mathcal{F}_{\lambda\mu}^r \mathcal{E}_r^\lambda - 2\nabla_\nu(\sigma^{\nu\beta} \mathcal{E}_{\mu\beta}) = 0,$$

where ∇_ν are covariant derivatives with respect to the Levi–Civita connection

$$K = dx^\lambda \otimes (\partial_\lambda + K_\lambda^\mu{}_\nu \dot{x}^\nu \partial_\mu), \quad K_\lambda^\mu{}_\nu = -\frac{1}{2} \sigma^{\nu\beta} (\sigma_{\lambda\beta\mu} + \sigma_{\mu\beta\lambda} - \sigma_{\beta\lambda\mu}).$$

7 Example II

Let us consider gauge theory of skew symmetric tensor fields. These are exterior forms on a base manifold X of degree more than one. We need not specify a gauge model, but refer to the topological BF theory [4]. This is a theory of two exterior forms A and B of form degree $|A| = \dim X - |B| - 1$. Another example is a gauge theory of an exterior form A in the presence of a background metric on X whose Lagrangian is similar to that of an electromagnetic field.

A generic gauge system of skew symmetric tensor fields is defined on the fiber bundle

$$Y = \bigwedge_X^p T^*X \oplus \bigwedge_X^q T^*X, \quad (7.1)$$

coordinated by $(x^\lambda, A_{\mu_1 \dots \mu_p}, B_{\nu_1 \dots \nu_q})$. The corresponding GDA is $\mathcal{O}_\infty^*[Y]$. There are the canonical p - and q -forms

$$A = \frac{1}{p!} A_{\mu_1 \dots \mu_p} dx^{\mu_1} \wedge \dots \wedge dx^{\mu_p} \in \mathcal{O}_\infty^{0,p}[Y], \quad B = \frac{1}{q!} B_{\nu_1 \dots \nu_q} dx^{\nu_1} \wedge \dots \wedge dx^{\nu_q} \in \mathcal{O}_\infty^{0,q}[Y] \quad (7.2)$$

on Y . A Lagrangian of the above-mentioned topological BF theory reads

$$L_{\text{BF}} = A \wedge d_H B, \quad p + q = n - 1. \quad (7.3)$$

A gauge symmetry of a generic gauge system of skew symmetric tensor fields, e.g., of the Lagrangian (7.3) is the following. Let us consider the fiber bundle

$$E = Y \times_X \bigwedge_X^{p-1} T^*X \times_X \bigwedge_X^{q-1} T^*X, \quad (7.4)$$

where

$$\bigwedge_X^{p-1} T^*X \times_X \bigwedge_X^{q-1} T^*X$$

is the fiber bundle of gauge parameters with coordinates $(x^\lambda, \varepsilon_{\mu_1 \dots \mu_{p-1}}, \xi_{\nu_1 \dots \nu_{q-1}})$. Let

$$\begin{aligned} \varepsilon &= \frac{1}{(p-1)!} \varepsilon_{\mu_1 \dots \mu_{p-1}} dx^{\mu_1} \wedge \dots \wedge dx^{\mu_{p-1}} \in \mathcal{O}_\infty^{0,p-1}[E], \\ \xi &= \frac{1}{(q-1)!} \xi_{\nu_1 \dots \nu_{q-1}} dx^{\nu_1} \wedge \dots \wedge dx^{\nu_{q-1}} \in \mathcal{O}_\infty^{0,q-1}[E] \end{aligned}$$

be canonical exterior forms like (7.2) on E (7.4). The above mentioned gauge symmetry is given by the generalized vector field

$$v = d_{\mu_1} \varepsilon_{\mu_2 \dots \mu_p} \frac{\partial}{\partial A_{\mu_1 \dots \mu_p}} + d_{\nu_1} \xi_{\nu_2 \dots \nu_q} \frac{\partial}{\partial B_{\nu_1 \dots \nu_q}}, \quad (7.5)$$

which acts on the exterior forms A and B (7.2) by the law

$$\mathbf{L}_{\vartheta_E} A = d_H \varepsilon, \quad \mathbf{L}_{\vartheta_E} B = d_H \xi. \quad (7.6)$$

In accordance with the formula (4.8), the corresponding Noether identity takes the form

$$-\varepsilon_{\mu_2 \dots \mu_p} d_{\mu_1} \mathcal{E}^{\mu_1 \dots \mu_p} = 0, \quad -\xi_{\nu_2 \dots \nu_q} d_{\nu_1} \mathcal{E}^{\nu_1 \dots \nu_q} = 0. \quad (7.7)$$

For instance, the equalities

$$\begin{aligned} \mathbf{L}_{\vartheta_E} L_{\text{BF}} &= (\mathbf{L}_{\vartheta_E} A) \wedge d_H B + A \wedge (\mathbf{L}_{\vartheta_E} d_H B) = d_H \varepsilon \wedge d_H B + A \wedge (\mathbf{L}_{\vartheta_E} d_H B) \\ &= d_H (\varepsilon \wedge d_H B) \end{aligned}$$

show that the generalized vector field (7.5) is a gauge symmetry of the Lagrangian L_{BF} (7.3). This Lagrangian provides the Euler–Lagrange equations

$$d_H A = 0, \quad d_H B = 0,$$

and the Noether identity (7.7) is brought into the form

$$d_H d_H A \equiv 0, \quad d_H d_H B \equiv 0.$$

It should be emphasized that the gauge symmetry (7.5) by no means exhausts all variational symmetries of the Lagrangian L_{BF} . Any generalized vector field v on Y such that $\mathbf{L}_{\vartheta} A$ is d_H -exact and $\mathbf{L}_{\vartheta} B$ is d_H -closed is a variational symmetry of this Lagrangian.

The gauge symmetry v (7.5) is reducible. Without a loss of generality, let us put $q \geq p$. Then v is q -stage reducible as follows. Let us consider vector bundles

$$\begin{aligned} E_k &= Y \times_X^{p-k-2} \bigwedge_X T^* X \times_X^{q-k-2} T^* X, \quad 0 \leq k < p-2, \\ E_k &= Y \times_X \mathbb{R} \times_X^{q-p} T^* X, \quad k = p-2, \\ E_k &= Y \times_X^{q-k-2} \bigwedge_X T^* X, \quad k > p-2, \end{aligned} \quad (7.8)$$

over Y provided with fiber coordinates

$$(\varepsilon_{\mu_1 \dots \mu_{p-k-2}}^k, \xi_{\nu_1 \dots \nu_{q-k-2}}^k), \quad (\alpha, \xi_{\nu_1 \dots \nu_{q-p}}^{p-2}), \quad (\xi_{\nu_1 \dots \nu_{q-k-2}}^k),$$

respectively. Then the differential operators

$$\begin{aligned} v_0 &= d_{\mu_1} \varepsilon_{\mu_2 \dots \mu_{p-1}}^0 \frac{\partial}{\partial \varepsilon_{\mu_1 \dots \mu_{p-1}}} + d_{\nu_1} \xi_{\nu_2 \dots \nu_{q-1}}^0 \frac{\partial}{\partial \xi_{\nu_1 \dots \nu_{q-1}}}, \\ v_k &= d_{\mu_1} \varepsilon_{\mu_2 \dots \mu_{p-k-1}}^k \frac{\partial}{\partial \varepsilon_{\mu_1 \dots \mu_{p-k-1}}^{k-1}} + d_{\nu_1} \xi_{\nu_2 \dots \nu_{q-k-1}}^k \frac{\partial}{\partial \xi_{\nu_1 \dots \nu_{q-k-1}}^{k-1}}, \quad 0 < k < p-2, \\ v_{p-2} &= d_{\mu} \alpha \frac{\partial}{\partial \varepsilon_{\mu}^{p-3}} + d_{\nu_1} \xi_{\nu_2 \dots \nu_{q-p+1}}^{p-2} \frac{\partial}{\partial \xi_{\nu_1 \dots \nu_{q-p+1}}^{p-3}}, \\ v_k &= d_{\nu_1} \xi_{\nu_2 \dots \nu_{q-k-1}}^k \frac{\partial}{\partial \xi_{\nu_1 \dots \nu_{q-k-1}}^{k-1}}, \quad k > p-2, \end{aligned}$$

satisfy the conditions of Definition 5.2.

Accordingly, there is a family of associated k -stage Noether operators Δ_k . Let the density duals $\overline{E}^* \rightarrow Y$, $\overline{E}_k^* \rightarrow Y$ of the vector bundles $E \rightarrow Y$ (7.4), $E_k \rightarrow Y$ (7.8) be provided with fiber coordinates

$$(\overline{\varepsilon}^{\mu_1 \dots \mu_{p-1}}, \overline{\xi}^{\nu_1 \dots \nu_{q-1}}), \quad (\overline{\varepsilon}_k^{\mu_1 \dots \mu_{p-k-2}}, \overline{\xi}_k^{\nu_1 \dots \nu_{q-k-2}}), \quad (\overline{\alpha}, \overline{\xi}_{p-2}^{\nu_1 \dots \nu_{q-p}}), \quad (\overline{\xi}_k^{\nu_1 \dots \nu_{q-k-2}}),$$

respectively. Then we have

$$\begin{aligned} \Delta_0 &= -[\varepsilon_{\mu_2 \dots \mu_{p-1}}^0 d_{\mu_1} \overline{\varepsilon}^{\mu_1 \dots \mu_{p-1}} + \xi_{\nu_2 \dots \nu_{q-1}}^0 d_{\nu_1} \overline{\xi}^{\nu_1 \dots \nu_{q-1}}] \omega, \\ \Delta_k &= -[\varepsilon_{\mu_2 \dots \mu_{p-k-1}}^k d_{\mu_1} \overline{\varepsilon}^{\mu_1 \dots \mu_{p-k-1}} + \xi_{\nu_2 \dots \nu_{q-k-1}}^k d_{\nu_1} \overline{\xi}^{\nu_1 \dots \nu_{q-k-1}}] \omega, \quad 0 < k < p-2, \\ \Delta_{p-2} &= -[\alpha d_{\mu} \overline{\varepsilon}_{p-3}^{\mu} + \xi_{\nu_2 \dots \nu_{q-p+1}}^{p-2} d_{\nu_1} \overline{\xi}_{p-3}^{\nu_1 \dots \nu_{q-p+1}}] \omega, \\ \Delta_k &= -\xi_{\nu_2 \dots \nu_{q-k-1}}^k d_{\nu_1} \overline{\xi}_k^{\nu_1 \dots \nu_{q-k-1}} \omega, \quad k > p-2. \end{aligned}$$

8 Appendix. Differential operators on dual fiber bundles

Given a fiber bundle $Y \rightarrow X$, let $E \rightarrow Y$, $Q \rightarrow Y$ be vector bundles coordinated by (x^λ, y^i, ξ^r) and (x^λ, y^i, q^a) , respectively. Let E^* , Q^* be their duals and $\overline{E}^*$, $\overline{Q}^*$ their density-duals (1.1) coordinated by (x^λ, y^i, ξ_r) , (x^λ, y^i, q_a) and $(x^\lambda, y^i, \overline{\xi}_r)$, $(x^\lambda, y^i, \overline{q}_a)$, respectively. Let v be a linear Q -valued differential operator on $E \rightarrow Y \rightarrow X$. It is represented by the function (2.10):

$$v = v^a q_a = \sum_{0 \leq |\Lambda| \leq m} v_r^{a, \Lambda} (x^\lambda, y_\Sigma^i) \xi_\Lambda^r q_a \in \mathcal{O}_\infty^0[E \times_Y Q^*], \quad 0 \leq |\Sigma| \leq m. \quad (8.1)$$

Let Δ be an m -order linear differential operator on the density-dual $\overline{Q}^* \rightarrow Y \rightarrow X$ of $Q \rightarrow Y$ with values in the density-dual $\overline{E}^*$ of E . It is represented by the function

$$\overline{\Delta} = \sum_{0 \leq |\Lambda| \leq m} \Delta_r^{a, \Lambda} (x^\lambda, y_\Sigma^i) \overline{q}_{\Lambda a} \overline{\xi}^r \in \mathcal{O}_\infty^0[(\overline{E}^*)^* \times_Y \overline{Q}^*], \quad 0 \leq |\Sigma| \leq m. \quad (8.2)$$

Let $J\omega$ be a volume form on X such that $\overline{\xi}_r = J\xi_r$ and $\xi^r = J\overline{\xi}^r$. Then the function (8.2) defines the density $\Delta = \overline{\Delta}J\omega$ which reads

$$\Delta = \Delta_r \xi^r \omega = \sum_{0 \leq |\Lambda| \leq m} \Delta_r^{a, \Lambda} (x^\lambda, y_\Sigma^i) \overline{q}_{\Lambda a} \xi^r \omega \in \mathcal{O}_\infty^{0, n}[E \times_Y \overline{Q}^*], \quad 0 \leq |\Sigma| \leq m. \quad (8.3)$$

Theorem 8.1. *Any linear Q -valued differential operator v (8.1) on $E \rightarrow Y \rightarrow X$ yields the linear $\overline{E}^*$ -valued differential operator*

$$\begin{aligned} \eta(v) &= \sum_{0 \leq |\Sigma| \leq m} (-1)^{|\Sigma|} d_\Sigma (v_r^{a, \Sigma} \overline{q}_a) \xi^r \omega = \sum_{0 \leq |\Lambda| \leq m} \eta(v)_r^{a, \Lambda} \overline{q}_{\Lambda a} \xi^r \omega, \\ \eta(v)_r^{a, \Lambda} &= \sum_{0 \leq |\Sigma| \leq m-|\Lambda|} (-1)^{|\Sigma+\Lambda|} C_{|\Sigma+\Lambda|}^{|\Sigma|} d_\Sigma (v_r^{a, \Sigma+\Lambda}), \end{aligned} \quad (8.4)$$

on $\overline{Q}^* \rightarrow Y \rightarrow X$. Conversely, any linear $\overline{E}^*$ -valued differential operator Δ (8.3) on $\overline{Q}^* \rightarrow Y \rightarrow X$ defines the linear Q -valued differential operator

$$\begin{aligned}\eta(\Delta) &= \sum_{0 \leq |\Sigma| \leq m} (-1)^{|\Sigma|} d_\Sigma(\Delta_r^{a,\Sigma} \xi^r) q_a = \sum_{0 \leq |\Lambda| \leq m} \eta(\Delta)_r^{a,\Lambda} \xi_\Lambda^r q_a, \\ \eta(\Delta)_r^{a,\Lambda} &= \sum_{0 \leq |\Sigma| \leq m-|\Lambda|} (-1)^{|\Sigma+\Lambda|} C_{|\Sigma+\Lambda|}^{|\Sigma|} d_\Sigma(\Delta_r^{a,\Sigma+\Lambda}),\end{aligned}\tag{8.5}$$

on $E \rightarrow Y \rightarrow X$. The relations (1.3) hold.

Proof. One must show that the differential operators given by the local coordinate expressions (8.4) and (8.5) are globally defined. The function v (8.1) yields the density

$$\overline{v} = \sum_{0 \leq |\Lambda| \leq m} v_r^{a,\Lambda} \xi_\Lambda^r \overline{q}_a \omega \in \mathcal{O}_\infty^{0,n}[E \times_Y Q^*].\tag{8.6}$$

Its Euler–Lagrange operator

$$\delta(\overline{v}) = \mathcal{E}_i dy^i \wedge \omega + \mathcal{E}_r d\xi^r \wedge \omega + \mathcal{E}^a d\overline{q}_a \wedge \omega$$

takes its values in the fiber bundle

$$V^*(E \times_Y Q^*) \otimes_{E \times_Y Q^*} \bigwedge^n T^*X,\tag{8.7}$$

where $V^*(E \times_Y Q^*)$ is the vertical cotangent bundle of the fiber bundle $E \times_Y Q^* \rightarrow X$. There is its canonical projection

$$\alpha_E : V^*(E \times_Y Q^*) \rightarrow V^*E \rightarrow V_Y^*E,\tag{8.8}$$

onto the vertical cotangent bundle V_Y^*E of $E \rightarrow Y$. Then we obtain a differential operator $(\alpha_E \circ \delta)(\overline{v})$ on $E \times_Y Q^*$ with values in the fiber bundle $V_Y^*E \otimes_E \bigwedge^n T^*X$. It reads

$$(\alpha_E \circ \delta)(\overline{v}) = \mathcal{E}_r \overline{d}\xi^r \otimes \omega = \sum_{0 \leq |\Lambda| \leq m} (-1)^{|\Lambda|} d_\Lambda(v_r^{a,\Lambda} \overline{q}_a) \overline{d}\xi^r \otimes \omega,$$

where $\{\overline{d}\xi^r\}$ is the fiber basis for $V_Y^*E \rightarrow E$ and the tensor product $\otimes$ is over $C^\infty(X)$. Due to the canonical isomorphism $V_Y^*E = E^* \times_Y E$, this operator defines the density (8.4). Conversely, the Euler–Lagrange operator of the density (8.3) takes its values in the fiber bundle (8.7) and reads

$$\delta(\Delta) = \mathcal{E}_i dy^i \wedge \omega + \mathcal{E}_r d\xi^r \wedge \omega + \mathcal{E}^a d\overline{q}_a \wedge \omega.\tag{8.9}$$

In order to repeat the above mentioned procedure, let us consider a volume form $J\omega$ on X and substitute $d\overline{q}_a \wedge \omega = Jdq_a \wedge \omega$ into the expression (8.9). Using the projection

$$\alpha_Q : V^*(E \times_Y Q^*) \rightarrow V_Y^*Q^*$$

like α_E (8.8) and the canonical isomorphism $V_Y^*Q^* = Q \times_Y Q^*$, we come to the density

$$\sum_{0 \leq |\Lambda| \leq m} (-1)^{|\Lambda|} d_\Lambda(\Delta_r^{a,\Lambda} \xi^r) q_a J \omega \in \mathcal{O}_\infty^{0,n}[E \times_Y Q^*]$$

and, hence, the function (8.5). The relations (1.3) result from the relation (2.19). $\square$

Relations (1.3) show that the intertwining operator η (8.4) – (8.5) provides a bijection between the sets $\text{Diff}(E, Q)$ and $\text{Diff}(\overline{Q}^*, \overline{E}^*)$ of differential operators (8.1) and (8.3).

Proposition 8.2. *Compositions of operators $v \circ v'$ and $\Delta' \circ \Delta$ obey the relations (1.4).*

Proof. It suffices to prove the first relation. Let $v \circ v' \in \text{Diff}(E', Q)$ be a composition of differential operators $v \in \text{Diff}(E, Q)$ and $v' \in \text{Diff}(E', E)$. Given fibred coordinates (ξ^r) on $E \rightarrow Y$, (e^p) on $E' \rightarrow Y$ and $(\overline{q}_a)$ on $\overline{Q}^* \rightarrow Y$, this composition defines the density (8.6)

$$\overline{v \circ v'} = \sum_\Lambda v_r^{a,\Lambda} d_\Lambda \left(\sum_\Sigma v_p^{r,\Sigma} \epsilon_\Sigma^p \right) \overline{q}_a \omega.$$

Following the relation (2.16), one can bring this density into the form

$$\sum_\Sigma v_p^{r,\Sigma} \epsilon_\Sigma^p \sum_\Lambda (-1)^{|\Lambda|} d_\Lambda(v_r^{a,\Lambda} \overline{q}_a) \omega + d_H \sigma = \sum_\Sigma v_p^{r,\Sigma} \epsilon_\Sigma^p \sum_\Lambda \eta(v)_r^{a,\Lambda} \overline{q}_{\Lambda a} \omega + d_H \sigma.$$

Its Euler–Lagrange operator projected to $V_Y^*E' \otimes_{E'} T^*X$ is

$$\sum_\Sigma (-1)^{|\Sigma|} d_\Sigma(v_p^{r,\Sigma} \sum_\Lambda \eta(v)_r^{a,\Lambda} \overline{q}_{\Lambda a}) \overline{d}e^p \otimes \omega = \sum_\Sigma \eta(v')_p^{r,\Sigma} d_\Sigma \left(\sum_\Lambda \eta(v)_r^{a,\Lambda} \overline{q}_{\Lambda a} \right) \overline{d}e^p \otimes \omega,$$

that leads to the desired composition $\eta(v') \circ \eta(v)$. $\square$

References

- [1] Anderson I 1992 Introduction to the variational bicomplex *Contemp. Math.* **132** 51-73
- [2] Barnich G, Brandt F and Henneaux M 2000 Local BRST cohomology in gauge theories *Phys. Rep.* **338** 439-569
- [3] Bashkirov D and Sardanashvily G 2005 On the BV quantization of gauge gravitation theory *Int. J. Geom. Methods Mod. Phys.* **2** No.2
- [4] Birmingham D, Blau M, Rakowski M and Thompson G 1991 Topological field theories *Phys. Rep.* **209** 129-340
- [5] Borowiec A, Ferraris M and Francaviglia M 2003 A covariant formalism of Chern–Simons gravity *J. Phys. A* **36** 2589-2598

- [6] Fatibene L and Francaviglia M 2003 *Natural and Gauge Natural Formalism for Classical Field Theories. A Geometric Perspective Including Spinors and Gauge Theories* (Dordrecht: Kluwer)
- [7] Fish M and Henneaux M 1990 Homological perturbation theory and algebraic structure of the antifield-antibracket formalism for gauge theories *Commun. Math. Phys.* **128** 627-640
- [8] Fulp R, Lada T and Stasheff J 2003 Noether variational Theorem II and the BV formalism *Rend. Circ. Mat. Palermo (2) Suppl.* No. 71 115-126
- [9] Giachetta G, Mangiarotti L and Sardanashvily G 2000 *Connections in Classical and Quantum Field Theory* (Singapore: World Scientific)
- [10] Giachetta G, Mangiarotti L and Sardanashvily G 2001 Cohomology of the infinite-order jet space and the inverse problem *J. Math. Phys.* **42** 4272-4282
- [11] Giachetta G, Mangiarotti L and Sardanashvily G 2003 Noether conservation laws in higher-dimensional Chern-Simons theory *Mod. Phys. Lett. A* **18** 2645-2651
- [12] Giachetta G, Mangiarotti L and Sardanashvily G 2005 Lagrangian supersymmetries depending on derivatives. Global analysis and cohomology. *Commun. Math. Phys.* (accepted) (*Preprint* hep-th/0407185)
- [13] Gomis J, París J and Samuel S 1995 Antibracket, antifields and gauge theory quantization *Phys. Rep* **295** 1-145
- [14] Gotay M 1991 A multisymplectic framework for classical field theory and the calculus of variations. In: *Mechanics, Analysis and Geometry: 200 Years after Lagrange* (Amsterdam: North Holland) 203-235.
- [15] Takens F 1979 A global version of the inverse problem of the calculus of variations *J. Diff. Geom.* **14** 543-562